

Papéis Avulsos de Zoologia

Museu de Zoologia da Universidade de São Paulo

Volume 46(7):67-71, 2006

www.scielo.br/paz

ISSN impresso: 0031-1049

ISSN on-line: 1807-0205

A NEW SPECIES OF THE GENUS *CRENILABIUM* (MOLLUSCA, HETEROBRANCHIA, ACTEONIDAE) FROM BRAZIL

LUIZ RICARDO L. SIMONE¹

ABSTRACT

Crenilabium birmani, a new species of acteonid is conchologically described from the southwestern and southern coast of Brazil. The new species is the first record of the genus for the South Western Atlantic and differ in details and proportions of the suture and aperture. Two variants of the suture were detected, some specimens have simple suture, while others possess a deep, scaled suture. Information about a possible specimen of *Crenilabium* figured by Abbott (1974) representing *Rictaxis punctocaelatus* is here discussed, and its holotype is figured.

KEYWORDS: *Crenilabium birmani* n. sp., Acteonidae, Brazil, deepwater, conchology.

INTRODUCTION

The acteonids are small gastropods that normally live in infratidal environment. The opisthobranch organization of their bodies contrasts with the heavy and well-developed shells. The genus *Crenilabium* Cossmann, 1889 [type species *Actaeon* (*C.*) *aciculatus* Cossmann, 1889:307 (pl. 8, fig. 30), OD, a Tertiary fossil from Europe] is sometimes considered subgenus of *Actaeon* Montfort, 1810 (e.g., Thiele, 1992:629). The genus encompasses species with elongated shell, few transversal lines at columellar margin of the shell aperture, and absence of fold at inner lip. The genus was never reported to Brazilian waters (Marcus, 1972; Rios, 1994).

In dredges off the southwestern and southern Brazilian coast, some shells belonging to *Crenilabium* were collected, revealing a new species formally described here.

MATERIAL AND METHODS

The specimens were obtained for study already dry, being constituted only by shells. They were examined in stereo-microscope and also in SEM in the "Laboratório de Microscopia Eletrônica", MZSP, some specimens were covered by gold.

Abbreviations of Institutions: MNRJ, Museu Nacional da Universidade Federal do Rio de Janeiro, Brazil; MZSP, Museu de Zoologia da Universidade de São Paulo, Brazil; USNM, National Museum of Natural History, Smithsonian Institution, Washington, D.C., USA.

Additional material for comparison with the species described here includes samples of the species *Rictaxis punctocaelatus* (Carpenter, 1864), beyond the holotype at USNM, another sample was examined: UNITES STATES OF AMERICA; California; San Luis Obispo County, Morro Bay, MNRJ HSL2443, 10 shells.

¹ Museu de Zoologia, Universidade de São Paulo, Caixa Postal 42494-970, CEP 04218-970, São Paulo, SP, Brasil. E-mail: lrsimone@usp.br

Systematics

Crenilabium birmani new species


(Figs. 1-12)

Types: Holotype MZSP 63161. Paratypes: BRAZIL; Rio de Janeiro; off São Tomé Cape, 21°42'S 40°15'W, 53 m depth, MZSP 63162, 1 shell (R.V. "W. Besnard" sta. Macaé 11, 23/ii/1992); 22°18'52.95"S 40°11'58.95"W, 190-200 m depth, MNRJ 10703, 9 shells (sta. P47; R.V. Astrogaroupa col. 18/xii/2004). Rio Grande do Sul;

off Rio Grande, 33°39'S 51°07'W, 200 m depth, MZSP 63163, 1 shell (R.V. "W. Besnard" sta. 1891; viii/1972).

Type locality: BRAZIL; São Paulo; off Ubatuba, 23°46'S 44°51'W, 55 m depth (R.V. "W. Besnard" sta. 1624; 16/xii/1971).

Diagnosis: Shell with spire as long as aperture. Suture simple or scaled, deep. Inner lip simple; inferior region of aperture rounded or weakly bended, lacking folds.




FIGURES 1-5. *Crenilabium birmani* shells in SEM: 1-3) holotype, 1) frontal view, length = 8.3 mm; 2) spire, frontal-slightly oblique view, scale = 0.5 mm; 3) detail of apex, profile, scale = 0.1 mm; 4-5) MZSP 63163 paratype; 4) detail of spire, frontal-slightly oblique view, scale = 0.5 mm; 5) frontal view, length = 7.6 mm.

Description

Shell (Figs. 1-9): Bulloid, antero-posteriorly elongated; maximum size 8 mm; approximately 2.5 longer than wide. Wall relatively thick. Color white or yellowish

white. Protoconch mammillated, of one whorl; separation with teleoconch marked by narrow orthocline furrow (Figs. 2, 3, 9). Teleoconch with up to 4.5 whorls, weakly convex, profile weakly rounded (almost straight). Suture marked by shallow, concavity and low



FIGURES 5-9. *Crenilabium birmani* shell of paratypes MNRJ 10703:6-8) frontal, right and dorsal views of larger specimen, length = 6.8 mm; 9) detail of apex in profile of a young specimen, scale = 0.25 mm; 10-11) *Riclaxia punctocaelatus* holotype USNM 14914 (courtesy of Ellen Strong, Smithsonian Institution); frontal and dorsal views; length = 4.9 mm.




FIGURE 12. Map of distribution of *Crenilabium birmani* (the symbols are slightly away from its actual place); star = specimen with simpler suture; lozenge = specimens with scaled suture.

and relatively wide, smooth furrow (Figs. 1, 2, 3); or deep furrow, with external edge away from superior surface of each whorl, similar to a scale turned superiorly (Figs. 4, 5, 6-9). Surface smooth, opaque, about 10 shallow, narrow spiral furrows, distributed rather irregularly; a narrow, spiral line located at certain distance from subsutural area. Last whorl with spiral furrows gradually becoming slightly deeper and close with each other towards umbilical area (Figs. 1, 5, 6, 7); about 15 in inferior half. Umbilicus absent. Aperture with about half of total length, antero-posteriorly elongated; superior end pointed, inferior (siphonal) region rounded (Figs. 1, 5, 6, 7). Inner lip concave; superior half to 2/3 convex, rounded, lacking callus; inferior half or 1/3 rounded, concave, with thick edge; between both regions of inner lip a strong notch in some specimens (Fig. 5). Outer lip simple, with cutting edge, subterminally thickened.

Measurements (respectively length and width in mm): Holotype: 8.3 by 3.1; MZSP 63162 (M11): 7.1 by 2.8; MZSP 63163: 7.6 by 2.9; MNRJ 10703 #1: 7.6 by 3.2; #2: 7.4 by 2.9.

Distribution (Fig. 12): From Rio de Janeiro (off São Thomé Cape) to north coast of São Paulo (off Ubatuba); also off Rio Grande, RS.

Habitat: Sandy bottoms; 53-200 m depth.

Material examined: Types.

Etymology: The specific epithet is in honor of the discoverer of the species, the physician and conchologist Adolpho Birman, from São Paulo, SP, Brazil.

DISCUSSION

There is variability in some characters of *Crenilabium birmani*. One of them is the dissimilar shape of the suture, in which some specimens have a simple and shallow furrow (Figs. 1, 2, 3), while it is deep, scaled in others (Figs. 4-9). The form of the aperture is also somewhat variable; the deepest portion of the inner lip is almost in middle level in some specimens (Fig. 1), while it is between middle and inferior thirds in others (Figs. 5, 7). The inferior or siphonal region of the aperture can be also different, it is rounded sometimes (Fig. 1, 6, 7), while it is somewhat squared in others (Fig. 5), having a clear angle between inner lip and siphonal edge. There is a coincidence between the two morphs described above, represented each one in the Figures 1 and 5 that induced the initial interpretation of two species. The morph represented in the Fig. 1 is found in shallower waters (53-55 m depth); while the other, represented in the Figs. 5 and 6, with scaled suture, is found deeper, at about 200 m. Despite a conservative approach is given here, as both morphs considered a single species, further studies can change this scenario. However, the suggestion is that both morphs are actually because of adaptations to different deepness. Related to the geographic distribution (Fig. 12), specimens with simpler suture (Fig. 1) are for moment only know from Rio de Janeiro to north São Paulo coast; in the mean time, specimens with scaled suture occur from Rio de Janeiro to the south coast of Rio Grande do Sul.

The placement in the genus is based mainly on the elongated shell shape, the relatively thick shell walls, and by the absence of folds at the inner lip. These characters are present in the species described herein that approach it from the type species *Crenilabium aciculatum* (see Zilch, 1959-1960:8, fig. 13). The fold in the inner lip is present in other elongated shelled genera of acteonids, such as *Tenuiacteon* Aldrich, 1921, and

Actaeonidea Gabb, 1873 (Zilch, 1959-1960:8-9), from which the presently described species can not belong, as it lacks this fold.

The described species can not be confused with any other *Crenilabium* species, however, it is similar to the Mediterranean and North Atlantic *C. exile* (Jeffreys, 1870) [= *Acteon nitidus* Verrill, 1882:540, pl. 58, fig. 21], from which *C. birmani* differs in having proportionally longer aperture, spire with deeper suture and last whorl proportionally longer. *Crenilabium birmani* additionally differs from the type species of the genus, *C. aciculatum*, by deeper suture, by shorter proportion of the aperture and by wider and less rounded shape of the anterior region of the aperture.

A specimen surprisingly similar to *Crenilabium birmani*, and to *Crenilabium exile*, is figured by Abbott (1974:312, fig. 3906) to represent the Californian *Rictaxis punctocaelatus* (Carpenter, 1864). On the other hand, the examination of the holotype of *R. punctocaelatus* (Figs. 10, 11) revealed completely different characters. Most subsequent pictures of that species never showed a different shape of the holotype [e.g., Marcus, 1972 (fig. 5); Rudman, 2001], which allow that Abbott's picture may be a misidentification of a *Crenilabium* species, possibly to the *C. exile* (species number 3908 of that catalogue). The species described here differ from that Abbott's specimen in having a proportional longer aperture and by less straight spire.

RESUMO

Crenilabium birmani, uma nova espécie de Acteonidae é descrita conquiliologicamente para a costa sudeste e sul do Brasil. A espécie nova é o primeiro registro do gênero para o sul do Atlântico Oeste e difere em detalhes e proporções da sutura e abertura. Duas variantes de sutura foram detectadas, alguns espécimes apresentam sutura simples, enquanto outros uma sutura funda e escalonada. Informações sobre um possível espécime de *Crenilabium* figurado por Abbott (1974), representando *Rictaxis punctocaelatus*, é discutido neste, com a ilustração do holótipo desta espécie.

PALAVRAS-CHAVE: *Crenilabium birmani* n. sp., Acteonidae, Brasil, águas profundas, conquiliologia.

ACKNOWLEDGMENTS

I special thank to Adolpho Birman by the specimens studied in this paper and subsequent donation to the MZUSP. To Dr. Alexandre Dias Pimenta, MNRJ, for the loan of MNRJ specimens and for comments on the manuscript. To Dr. Ellen Strong, Smithsonian Institution, Washington, D.C. by the photo of *Rictaxis punctocaelatus* holotype. To Lara Guimarães, Laboratório de Microscopia Eletrônica, MZSP, in helping the SEM examination of the specimens. This study is partially supported by FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo) projects, numbers 04/00309-2; 04/10793-9.

REFERENCES

- Abbott, R.T. 1974. *American Seashells*. Second edition. Van Nostrand Reinhold Company, New York, 663 p. + 240 pls.
- Carpenter, P.P. 1864. Supplementary report on the present state of our knowledge with regard to the mollusca of North America. *Report of the British Association for the Advancement of Science*, 1863:517-686.
- Cossmann, M. 1889. Catalogue Illustré des coquilles fossiles de l'Éocène des Environs de Paris. (4ème fascicule). *Annales de la Société Royale Malacologique de Belgique*, 24:3-385 + pl. I-XII.
- Marcus, E.B.R. 1972. On Some Acteonidae (Gastropoda, Opisthobranchia). *Papéis Avulsos de Zoologia (São Paulo)*, 25(19):167-188.
- Rios, E.C. 1994. *Seashells of Brazil*. Second edition. Fundação Universidade do Rio Grande, Rio Grande, 368 p. + 113 pls.
- Rudman, W.B. 2001. *Rictaxis punctocaelatus* (Carpenter, 1864). In: *Sea Slug Forum*. Australian Museum, Sydney. Available at: <http://www.seaslugforum.net/factsheet.cfm?base=rictpunc>
- Thiele, J. 1992. Handbook of Systematic Malacology. Part 2 (Gastropoda: Opisthobranchia and Pulmonata). In: Bieler, R & Mikkelsen, P. [translators]. *Smithsonian Institution Libraries*. The National Science Foundation, Washington, p. 627-1189.
- Verrill, A. E. 1882. Catalogue of marine Mollusca added to the fauna of the New England region, during the past ten years. *Transactions of the Connecticut Academy of Arts and Sciences*, 5:451-587 + pls. 42-44, 57-58.
- Zilch, A. 1959-1960. Euthyneura. In: W. Wenz. *Gastropoda*. Handbuch der Paläozoologie. Berlin. Band 6(2):xii + 834 p.

Recebido em: 20.04.2006

Aceito em: 23.08.2006

EDITORIAL COMMITTEE

Editor-in-Chief: Hussam Zaher, Serviço de Vertebrados, Museu de Zoologia, Universidade de São Paulo, Caixa Postal 42.494, CEP 04218-970, São Paulo, SP, Brasil. E-mail: hzaher@ib.usp.br.

Associate Editors: Carlos José Einicker Lamas (Universidade de São Paulo, Brasil); Antonio Carlos Marques (Universidade de São Paulo, Brasil); Mário César Cardoso de Pinna (Universidade de São Paulo, Brasil); Sergio Antonio Vanin (Universidade de São Paulo, Brasil).

Editorial Board: Aziz Nacib Ab'Saber (Universidade de São Paulo, Brasil); Rüdiger Bieler (Field Museum of Natural History, U.S.A.); Walter Antonio Pereira Boeger (Universidade Federal do Paraná, Brasil); Carlos Roberto Ferreira Brandão (Universidade de São Paulo, Brasil); James M. Carpenter (American Museum of Natural History, U.S.A.); Ricardo Macedo Corrêa e Castro (Universidade de São Paulo, Brasil); Darrel R. Frost (American Museum of Natural History, U.S.A.); William Ronald Heyer (National Museum of Natural

History, U.S.A.); Ralph W. Holzenthal (University of Minnesota, U.S.A.); Adriano Brillhante Kury (Museu Nacional, Rio de Janeiro, Brasil); John G. Maisey (American Museum of Natural History, U.S.A.); Naércio Aquino Menezes (Universidade de São Paulo, Brasil); Christian de Muizon (Muséum National d'Histoire Naturelle, Paris, France); Gerardo Lamas Müller (Museo de Historia Natural "Javier Prado", Lima, Peru); Nelson Papavero (Universidade de São Paulo, Brasil); James L. Patton (University of California, Berkeley, U.S.A.); Richard O. Prum (University of Kansas, U.S.A.); Marcos André Raposo (Museu Nacional, Rio de Janeiro, Brasil); Olivier Rieppel (Field Museum of Natural History, U.S.A.); Miguel Trefaut Urbano Rodrigues (Universidade de São Paulo, Brasil); Randall T. Schuh (American Museum of Natural History, U.S.A.); Ubirajara Ribeiro Martins de Souza (Universidade de São Paulo, Brasil); Marcos Tavares (Universidade de São Paulo, Brasil); Paulo Emilio Vanzolini (Universidade de São Paulo, Brasil); Richard P. Vari (National Museum of Natural History, U.S.A.); Mario de Vivo (Universidade de São Paulo, Brasil); Paulo Secchin Young (Museu Nacional, Rio de Janeiro, Brasil).

INSTRUCTIONS TO AUTHORS (MAY 2002)

General Information: *Papéis Avulsos de Zoologia* covers primarily the fields of Zoology, publishing original contributions in systematics, paleontology, evolutionary biology, ecology, taxonomy, anatomy, behavior, functional morphology, molecular biology, ontogeny, faunistic studies, and biogeography. *Papéis Avulsos de Zoologia* also encourages submission of theoretical and empirical studies that explore principles and methods of systematics.

Lists and catalogs should be submitted to the Arquivos de Zoologia.

All contributions must follow the International Code of Zoological Nomenclature. Relevant specimens should be properly curated and deposited in a recognized public or private, non-profit institution. Tissue samples should be referred to their voucher specimens and all nucleotide sequence data (aligned as well as unaligned) should be submitted to GenBank (<http://www.ncbi.nih.gov/Genbank/>) or EMBL (<http://www.ebi.ac.uk/>).

Peer Review: All submissions to *Papéis Avulsos de Zoologia* are subject to review by at least two referees and the Editor-in-Chief. Three legible copies (including photocopies of original illustrations) and original illustrations must be submitted; all authors will be notified of submission date. Authors may suggest potential reviewers. Communications regarding acceptance or rejection of manuscripts are made through correspondence with the first or corresponding author only. Once a manuscript is accepted providing changes suggested by the referees, the author is requested to return a revised version incorporating those changes (or a detailed explanation of why reviewer's suggestions were not followed) within four weeks upon receiving the communication by the editor. Revised manuscripts must be submitted as both hard copy and electronic file (3.5" disk, Zip Drive, or CD Rom with text in Microsoft Word format). Figures and graphics should be sent separately (".jpg", ".tif", ".xls", ".cdr").

Proofs: Page-proofs with the revised version will be sent to the first or corresponding author. Page-proofs must be returned to the editor in two weeks, preferentially within 48 hours. Failure to return the proof promptly may be interpreted as approval with no changes and/or may delay publication. Only necessary corrections in proof will be permitted. Once page proof is sent to the author, further alterations and/or significant additions of text are permitted only at the author's expense or in the form of a brief appendix ("note added in proof").

Submission of Manuscripts: Manuscripts should be sent to the Editor-in-Chief (H. Zaher, Museu de Zoologia da USP, Caixa Postal 42.494, CEP 04218-970, São Paulo, SP, Brasil). Manuscripts are considered on the understanding that they have not been published or will not appear elsewhere in substantially the same or abbreviated form. The criteria for acceptance of articles are: quality and relevance of research, clarity of text, and compliance with the guidelines for manuscript preparation.

Manuscripts should be written preferentially in English, but texts in Portuguese or Spanish will also be considered. Studies with a broad coverage are encouraged to be submitted in English. All manuscripts should include an abstract in Portuguese and English regardless of the original language.

Authors are requested to pay attention to the instructions concerning the preparation of the manuscripts. Close adherence to the guidelines will expedite processing of the manuscript, whereas manuscripts deviating from the required form will be returned for revision prior to review.

Manuscript Form: Manuscripts should not exceed 100 pages of double-spaced typescript on 21 by 29.7 cm (A4 format) or 21.5 by 28 cm (letter format) paper, with wide margins. The pages of the manuscript should be numbered consecutively.

The text of articles should be arranged in the following order: Title Page, Abstracts, Body of Text, Literature Cited, Tables, Appendices, and Figure Captions. Each of these sections should begin on a new page. All typescript pages must be double-spaced.

- (1) **Title Page:** This should include the title, author(s) name(s), institutions, and keywords in English as well as in the language of the manuscript, and a short

running title in the language of the manuscript. The title should be concise and, where appropriate, should include mention of families and/or higher taxa. Names of new taxa should not be included in titles.

- (2) **Abstract:** All papers should have an abstract in English and another in Portuguese, regardless of the original language. The abstract is of great importance as it may be reproduced elsewhere. It should be in a form intelligible if published alone and should summarize the main facts, ideas, and conclusions of the article. Telegraphic abstracts are strongly discouraged. Include all new taxonomic names for referencing purposes. Abbreviations should be avoided. It should not include references. Abstracts should not exceed 350 words.
- (3) **Body of Text:** The main body of the text should include the following sections: Introduction, Materials and Methods, Results, Discussion, and Acknowledgments at end. Primary headings in the text should be in capital letters and centered; the following text should begin on the next line, indented. Secondary headings should be in capital and lowercase letters and flush left; the following text should begin on the next line, indented. Tertiary headings should be in capital and lower case letters, in italics and indented; the following text should be on the same line and separated from the heading by a hyphen.
- (4) **Literature Cited:** Citations in the text should be given as: Silva (1998)..., Silva (1998:14-20)..., Silva (1998: figs. 1, 2)..., Silva (1998a, b)..., Silva & Oliveira (1998)..., (Silva, 1998)..., (Rangel, 1890; Silva & Oliveira, 1998a, b; Adams, 2000)..., (Silva, pers. comm.)..., (Silva *et al.*, 1998), the latter when the paper has three or more authors. The reference need not be cited when author and date are given only as authority for a taxonomic name. The literature section should be arranged strictly alphabetically and given in the following format:

Journal Article – Silva, H.R.; Oliveira, H. & Rangel, S. Year. Article title. Journal name, 00:000-000. Names of journals must be spelled out in full.

Books – Silva, H.R. Year. Book title. Publisher, Place.

Articles in Books – Silva, H.R. Year. Article title. In: Oliveira, H. & Rangel, S. (Eds.), Book title. Publisher, Place, p.000-000.

Articles in Larger Works – Silva, H.R. Year. Article title. In: H. Oliveira & S. Rangel (Eds.), Title of Larger Work. Serial Publication. Publisher, Place, p.000-000.

Dissertations and Theses – Silva, H.R. Year. Dissertation title. Ph.D. Dissertation, University, Place.

Electronic Publications – Silva, H.R. Year. Article title. Available at: <http://www.mz.usp.br>.

Tables: All tables must be numbered in the same sequence in which they appear in the text. Authors are encouraged to indicate where the tables should be placed in the text. They should be comprehensible without reference to the text. Tables should be formatted with horizontal, not vertical, rules. In the text, tables should be referred as Table 1, Tables 2 and 3, Tables 2-6. Use "TABLE" in the table heading.

Illustrations: Figures should be numbered consecutively, in the same sequence they appear in the text. Separate illustrations of a composite figure should be identified by capital letters and referred in the text as so (fig. 1A). Where possible, letters should be placed in the lower right corner of each illustration of a composite figure. Hand-written lettering on illustrations is unacceptable. Illustrations should be mounted on stout, white cardboard. Figures should be mounted in order to minimize blank areas between separate illustrations. High quality color or black and white photographs, and computer generated figures are preferable. Authors are encouraged to indicate where the figures should be placed in the text. Use "(Fig(s).)" and "Figure(s)" for referring to figures in the text, but "FIGURE(S)" in the figure captions and "(fig(s).)" when referring to figures in another paper.

For other details of manuscript preparation of format, consult the CBE Style Manual, available from the Council of Science Editors (<http://www.councilscienceeditors.org/publications/style.cfm>).

Papéis Avulsos de Zoologia and *Arquivos de Zoologia* are publications of the Museu de Zoologia da Universidade de São Paulo (www.mz.usp.br).